

Ear wax in your child: what to do with the goo



Babies are gooey. Spew tends to dribble out of every orifice and the ear is no exception.

Devin's mother tipped her four month old baby's head sideways in the office the other day and asked me what to do about the oily, yellow wax smeared around the opening of his ear canal. Despite the copious amount of wax on the outside, Devin's ear canals were clear. "But the wax is simply disgusting," said Devin's mom, "Can I clean his ears? "

If you can get the wax with a wash cloth, it's fair game. Otherwise, leave it alone. It doesn't matter if you use a wash cloth or cotton swab. The special shaped cotton swabs with the safety tips are unnecessary. Rest assured, you will not go too deeply into the ear canal if you only scrape off what is visible.

Now suppose your pediatrician does say the wax should be removed. Place an over-the-counter solution such as Debrox in the ears (children and adults can use the same formulation) – three to four drops one or two times a day (during sleep is easiest for babies and toddlers) for a few days. The solution

softens wax. For maintenance, mineral oil and olive oil are favorite remedies. Place one drop daily in ears. In the office some pediatricians can use a water irrigation system (like a water squirter in your ear) to wash out the wax. The worst side effect is that the child's shirt sometimes gets wet. Irrigation is a very effective for removing wax in a school-aged or teenaged child who complains of difficulty hearing.

Some say wax evolved to help keep bugs and other debris from reaching deep into our ear canals. Case in point: one of my least favorite memories during residency is of picking out pieces of a cockroach entrapped in a child's earwax!

Keep in mind the amount of wax you see on the outside of the ear is not indicative of the actual amount inside the ear canal. Chances are, the wax is not hard and does not block the ear drum. Even if there is a large amount of wax, it is unlikely to greatly affect a baby's hearing unless the wax is stuck against the ear drum. Equally normal is that some babies and children don't seem to produce any ear wax. If you are concerned about your child's ear wax or about her hearing, have your pediatrician take a peek with a light.

If you find you are constantly cleaning your kid's waxy ears, take heart. At least there won't be any roaches "bugging" them.

Naline Lai, MD and Julie Kardos, MD

©2011, 2016 Two Peds in a Pod®

PS: Medical vocabulary FYI: light used to look into ears= otoscope. Medical term for ear wax= cerumen.

We give thanks, 2016

Nearly seven years ago, on the swimming pool bleachers at the local Y, I happened to sit next to Lexi Logan. Above the echoing din of kids splashing, I discovered that although she was trained as a painter, Lexi was interested in branching out into photography. Coincidentally, Dr Kardos and I were interested in branching medicine out into a new media called the internet and were dismayed at the lack of publicly available photos to accompany our blog posts. Lexi and I intersected in the right place at the right time. Since that chance meeting, Lexi has generously shared dozens of photos with Two Peds in a Pod.

The woman in the photo below, between your Two Peds (Dr. Kardos with the curly hair, Dr. Lai with the straight hair), is our photographer extraordinaire, Lexi Logan. Her work, which you can check out at www.lexilogan.com, speaks for itself. Local peeps may want to contact her to take their own family photos.

This Thanksgiving we say thanks to all those parents we've ever sat next to on bleachers. All the kid-related information we have learned, from navigating chorus uniforms, bus stop times, best teachers, fun summer camps, and even starting up blogs, has been invaluable.

In particular- thank you, Lexi!

We wish all of our readers a very healthy and happy Thanksgiving,

Dr. Naline Lai with Dr. Julie Kardos

©2016 Two Peds in a Pod®



**Extra school services offered
to your child? Just say yes!**



When I was in first and second grade, I took “special gym.”

I attended a public school in a small New Jersey town. The school building was about 100 years old, and the “special services” that my school offered were speech, reading help, and special gym.

I remember being THRILLED that I was selected to take special gym, because instead of just *one* day a week of bouncing balls and running races and turning somersaults during the school day, I got to go *twice* a week. I remember how upbeat and energetic the gym teacher was, and how much fun she made these exercises. I do not recall such words as “physical therapy” or “occupational therapy.” In fact, I did not realize the true point of the extra gym days until many years later, when I was in college and reminiscing about elementary school and caught myself mid-sentence:

“Well, when I was in first grade, I took special gym... hey... WAIT a MINUTE...!”

That’s when I realized that I had been flagged with a coordination challenge. Unbeknownst to me, in school I went to

physical therapy weekly.

Now that first-quarter parent teacher conferences are over, you may be surprised that your child has been offered special services by the school. Teachers spend hours a day with our kids and are experts in the age group that they teach. Not all kids are good at learning all subjects and not all are equally sociable or equally physically adept. When teachers ask a parent's permission to supply extra help, parents should not take this request as an affront or attack on their parenting. Rather, it is an opportunity to help kids succeed.

I was never suspicious about my inclusion in special gym. No one made fun of me for being in the class, and in fact many were jealous. Kids in early grades may be aware that some of their classmates come and go during the day, but they do not distinguish between kids pulled out for a gifted program from kids pulled out for remedial education. As an adult, I appreciate that my teachers made me feel good about being included in the special gym club.

I have a magnet on my car now that says, simply, "13.1," which is the number of miles that I ran to complete the Trenton Half-Marathon this past October. Special gym did not hold me back—it propelled me forward. I had no idea that my participation in special gym was emotionally charged for my mom until after I called my dad to tell him my race time (2 hours, 11.5 minutes). Only then did he tell me how crushed my mom had been about my inclusion in special gym. I am grateful that she hid that from me.

My message: Let your kids get extra help in school, allow them to be pulled out of a class they are failing and placed into an environment where they can learn and overcome challenges. Allow yourself to mourn the loss of the child you may have pictured. But know this: young children do not have enough life experience to independently think of themselves as failures in the early school years. They look to adults who

are important to them for how to respond to challenges and frustration. Encourage them with the positive message that they will receive extra attention and extra time to work at reading or math or physical skills or speech skills. Who knows? They may become the kid who applies to medical school or runs a marathon (or a half-marathon) someday.

Julie Kardos, MD

©2016 Two Peds in a Pod®

Sore throat relief



The giraffe always felt his sore throat lasted longer than everyone else's sore throat.

Many times parents bring their children with a sore throat

to our office to “check if it’s strep.” Some are disappointed to find out that their child does NOT have strep. Moms and Dads lament, “But what can I do for him if he can’t have an antibiotic? At least strep is treatable.”

Take heart. Strep or no strep, there are many **ways to soothe your child’s sore throat:**

- **Give pain medication** such as acetaminophen (brand name Tylenol) or ibuprofen (brand names Advil or Motrin). Do not withhold pain medicine before you bring her in to see her pediatrician. Too many times we hear parents say, “We wanted you to see how much pain she is in.” No need for this! Pediatricians are all in favor of treating pain as quickly and effectively as possible. Pain medicine will not interfere with physical exam findings nor will it interfere with strep test results.
- **Give lots to drink.** Some kids prefer very cold beverages, others like warm tea or milk. Avoid citrus juices since they sometimes sting sore throats. Frozen Slurpies or milkshakes, on the other hand, feel great on sore throats. Tell your child that the first three sips of a very cold drink may hurt, but then the liquid will start to soothe the throat. Watch for signs of dehydration including dry lips and mouth, no tears on crying, urination less than every six hours, and lethargy.
- **Provide soft foods** if your child is hungry. For example, noodles feel better than a hamburger on a sore throat. And ice-cream or sherbet therapy is effective as well.
- **Try honey** (if your child is older than one year) – one to two teaspoons three times a day. Not only can it soothe a sore throat but also it might quiet the cough that often accompanies a sore throat virus. Give it alone or mix it into milk or tea.
- **Kids older than three years** who don’t choke easily can suck on lozenges containing pectin or menthol for relief. Warning: kids sucking on lozenges may dupe

themselves into thinking they are hydrating themselves. They still need to drink to stay hydrated.

- **Salt water gargles** are an age-old remedy. Mix 1 teaspoon of salt in 6 ounces of warm water and have your kid gargle three times a day.
- **Magic mouthwash:** For those older than 2 years of age, mix 1/2 teaspoon of liquid diphenhydramine (brand name Benadryl 12.5mg/5ml) with 1/2 teaspoon of Maalox Advanced Regular Strength Liquid (ingredients: aluminum hydroxide, magnesium hydroxide 200 mg, and simethicone) and give a couple times a day to coat the back of the throat prior to meals. The Maalox coats the throat and the benedryl acts as a weak topical anesthetic (pain reliever). **Do not** use the Maalox formulation which contains bismuth subsalicylate because bismuth subsalicylate is an aspirin derivative, and aspirin is linked to [Reye's syndrome](#).
- For kids three years and older, **try throat sprays** containing phenol (brand name Baker's P&S and Chloraseptic® Spray for Kids). Use as directed.

Strep throat typically does **not** cause a bad cough, profuse runny nose, ulcers in the throat, or laryngitis. If your child has these other symptoms in addition to her sore throat, you can be fairly sure that she does NOT have strep. For a better understanding of strep throat, see our updated post on this topic.

The following are each a very important sign that a child with a **sore throat needs to see a doctor** for further evaluation:

1-can't swallow (kids might even spit out their own saliva)

2-can't open his mouth fully

3- hurts so much that the pain is not alleviated with the above measures in this post

4- presence of fever 101F or higher for more than 3-4 days

5-is accompanied by a new rash

Please also see our prior post on [how to tell if you need to call your child's doctor for illness.](#)

Julie Kardos, MD and Naline Lai, MD

©2016, 2015, 2012 Two Peds in a Pod®

Stop a bully: talk, walk, squawk, and support



photo by Lexi Logan

Can you identify your child in any of these scenarios?

-Your second grader comes off the school bus crying because another student was teasing him the entire ride home about his new glasses.

-Your fifth grader was punched on the school yard by a sixth grader and none of the playground teachers saw it happen. Your child's friend shoved the older child off your child before more damage was done.

-Your ninth grader keeps missing the school bus so you have to drive her to school. When she comes home from school she uses the bathroom immediately. You find out she avoids the bus and the school bathroom because kids make fun of her in both places.

Whatever your child's age, when you realize he or she is being bullied you will be outraged. In fact you might be tempted to retaliate against the bully yourself. However, here are more appropriate ways to help your child.

Bullying should never be tolerated. Teach your child how to directly deal with a bully, but be quick to talk also to the adult supervising your child when the bullying occurs. Your child should always feel safe in school, day camp, on a sports team, or any other adult-supervised activity.

Bullies are always in a position of power over their victims; either they are physically larger, older, or more socially popular. Teach your child first to try a strong verbal response (**talk**) such as "STOP talking to me that way!" or "Don't DO that to me!" Speaking strongly and looking the bully in the eye may take away some of the bully's power as well as attract attention of nearby peers or adults who can help your child.

Teach your child to **walk** away from a fight. Tell him to keep on walking toward a teacher, a classroom, a peer, or anyone else who can offer safety from a bully. Train him to breathe deeply/ignore/de-escalate situations to diffuse a bully's anger.

Have your child tell a teacher, camp counselor, coach, or other supervising adult about the abuse (**squawk**) as soon as it occurs. Always encourage your children to talk to you as well. Remember at home to ask your child questions such as "How is school," "How are your friends," "Do you know any kids who are being bullied?," and "Are YOU being bullied?" Dr. Lai always advises her patients to tell as many different adults as possible if he is not feeling safe. Even if one adult is unsure of how to help, sooner or later some one will.

If your child says he is angry at a friend or a classmate, be sure to ask questions that encourage your child to elaborate, such as "Oh, what happened?" or "Did something happen between you?" Listen carefully to his response. He may be taking out his anger at a bully on his own friends. This response is in retaliation for his friend's failure to protect him from a bully. Also, is your child becoming more reluctant to attend school, "missing" the bus more often and thus requiring a ride, or acting angry or sad more often? Kids who are victims of bullying can act like this.

In school, once you are aware that your child is a victim, talk not only to your child about how she should handle a bully but also alert your child's teacher and/or school principal about the situation (**support**). You should tell them in your child's words what happened, what was said, and be clear that you are asking for more supervision so that the bully has less access to your child. Ask for more supervision during times when there is usually less adult presence such as in the lunchroom or on the schoolyard. Your school may already have a "no bullying" policy. Often, the aggressor gets the heavier consequence in the event of a conflict. Again, children have a right to feel safe in school.

Restore your child's self-confidence. Bullies pick on kids who are smaller and weaker than they are, physically as well as psychologically. So your child has more positive experiences with kids who do not bully, encourage your child to invite

friends over to your home or host a fun group activity (kickball game in your backyard, show a movie/supply popcorn, etc.). Do family activities and show your child that you enjoy spending time with him. Enroll your child in activities that increase his self esteem such as karate, sports, or music lessons. A child who feels good about himself "walks taller" and is less likely to attract a bully

As a parent, you might read this post and think, "Yes, but I'd rather just teach my child to take revenge." Unfortunately, escalating the situation only breeds anger and in fact may get your child into trouble. Rather than "hate" the bully, help your child see that a bully deep down feels insecure. A bully resorts to making himself feel better by making others feel bad. Teach your child to pity the bully. With your guidance, your child will project self-confidence and a bully will never, ever, be able to touch him.

While the topic of cyper-bullying could occupy an entire separate post, we just want to alert you to the power that social media has over our kids as well. Ask your kids and teens directly about bullying that occurs on-line just as you would ask about bullying at school. Virtual bullying, unfortunately, is just as potentially harmful as in-person bullying, and is a known risk factor for teen suicide. Remind your children how important it is to refrain from revenge: better to disengage from social media than to respond to on-line bullying because your child will leave a permanent footprint on their on-line presence. Lay down the general rule of never posting anything negative (even a simple "dislike") online.

Help your child talk, walk, squawk and seek support. All kids deserve to feel secure in themselves and in the world around them.

Additional resources:

The American Academy of Pediatrics

Stopbullying.gov–Bully prevention site managed by U.S. Department of Health and Human Services

Cyberbullying Research Center–an organization dedicating to providing up to date information on cyberbullying

Teaching tolerance– a site where parents and educators can learn ways to foster tolerance

Julie Kardos, MD and Naline Lai, MD

© 2016, 2010 Two Peds in a Pod®

How to raise an optimist (even in an election year)



In this time of United States “election stress,” we bring back guest bloggers psychologist Dr. Gage and pediatrician Dr. Penaflor’s post on how to build optimism in your children.

Recently, my daughter’s friend announced before a race, “I’m just not going to try my best.”

Why would a child give up before even starting? Why such pessimism?

It turns out that her friend’s mother would say after every

race, “You just didn’t meet your potential. Did you at least beat Sarah (a fellow competitor)?”

This scenario illustrates how a parent who constantly gives negative responses can build pessimism in a child.

Why is optimism important?

An optimistic child is strong, enterprising, and resilient. He or she does not wait passively for good things to happen to him or her. The optimist consciously plans, works hard to make things happen, and persists through challenges.

Research shows important benefits:

- A healthier heart and a greater ability to fight infections and survive disease
- Better response to stress
- Less likelihood to develop anxiety and depression
- More success in school, sports, social and recreational activities
- Greater accomplishments in life

How do I begin?

Does your child tend toward optimism or pessimism? Is the glass half empty or half full? Which would your child say, “It doesn’t matter... I won’t get it right anyway,” or “I did my best... I’ll get it next time”? Optimism is a learned skill that you can teach your child at home.

Here are some important tips.

Model positive behaviors and attitudes:

“This is tough, but I can do it!”

“I will find that lost pair of socks!”

Create an environment that **fosters love and trust**.

When children have a sense of security and trust at home, they view the world as a positive place to explore and try new things.

Encourage your child to view life in a positive way and to rise above negativity.

For example, one of our favorite techniques is “**Rise up! Don’t dwell on it.**” If someone did or said something hurtful to your child, teach your child to pause. Have her ask herself “How important is it? Will it matter in 5 minutes, 5 months, or in 5 years?” Think of the big picture.

Another is to approach mistakes calmly. Say “Oops!” and move on.

Validate your child’s feelings of disappointment or sadness, but teach your child that failures and mistakes are opportunities to learn and do something different and better.

After all, in life “Sometimes you win, sometimes you learn.”

Patricia Gage, PhD, NCSP and Gina Penafior, MD, FAAP

©2016, 2014 Two Peds in a Pod®

***Dr. Patricia Gage** runs Brain Smart Academics, her own private practice as a school psychologist in Stuart, Florida, and has taken the lead in many charitable organizations that help promote children’s social/emotional wellness and women’s health.*

***Dr. Gina Penafior**, mother of a school-aged child, tween and teen, is a primary care pediatrician in South Florida with a background in emergency and hospital medicine. She and Pat have combined their knowledge and experiences to create a Hang-In-There educational card series. Their mission is to help busy moms and dads lead a more rewarding (and less stressful!) parenting experience.*

To learn more, please visit their website at www.HangNthere.com or Facebook page, or e-mail them at busymoms@HangNthere.com.

Potty training: the advanced course



Just when you thought your child graduated from potty training ...there's more

Believe it or not, this post is for parents of kids who are already potty trained. Because sometimes even kids who are potty trained will refuse to use the toilet.

Changes and transitions, such as the start of a new

school year, can trigger regression in kids who have been potty trained for years. Now that your children have been back to school for a few weeks take the quiz:

-Does your child come home from school and make **a mad dash for the bathroom** before he even gets his shoes off or asks you for a snack?

-Does your child stop eating and complain of **belly pain after two bites of dinner?**

-Does your child's **poop routinely clog your toilet?**

-Is your child's **underwear sometimes damp?**

-Have you noticed **skid marks** (small streaks or smears of poop) on your child's underwear?

If you answer "yes" to ANY of these questions, then read on.

Advanced Potty Training

The main problem many kids encounter is that once school starts, they have a lot of new distractions and can't be bothered to pee or poop. First, the morning routine might be more rushed. In school, the teacher is teaching. Your kid is interacting with other kids. The school's bathroom is foreign, and may even have an auto-flush toilet: scary for the newly-trained. And unlike your newly potty-trained toddler who often finds it thrilling to try out every public restroom he sees, kids in school may feel more self-conscious, and not as adventurous, about visiting new bathrooms.

School bathrooms can be smelly, loud, and even places where kids bully each other. Some kids develop an aversion to using the school bathroom. These kids hold their pee and poop all day long until they get home, then run into the house and make a mad dash for the bathroom.

A child's internal debate

For a kid who becomes overstimulated at school or who develops an aversion to the school bathroom, his bladder and brain have a conversation that goes like this:

Bladder: I am full.

Brain: Hold it, I don't want to use the bathroom right now.

Bladder: But I REALLY have to pee.

Brain: Tough luck, Bladder, just wait till we get home.

Bladder: But I have no more room for pee!

Brain: Deal with it, Bladder!

So, the bladder has two choices:

1-Bladder overflows, at least enough to relieve a bit of pressure. This causes damp underwear. (For other reasons click [here](#) to review our post on damp underwear.)

OR

2-Bladder distends to accommodate more urine and confuses the nerves that supply sensation to the bladder. Kids lose the ability to tell if they have to urinate, which leads to full bladder-emptying accidents, and can lead to urinary tract infections.

A similar discussion can occur between your kid's brain and his rectum:

Rectum: I am full of poop, Brain. Take us to the bathroom.

Brain: In school? Are you kidding? I am enjoying this game the class is playing/I am embarrassed to poop in school/I am afraid of the school bathroom. HOLD IT!

Rectum: OK, but you're not going to like how this comes out in the end.

The rectum also has two choices: release just a bit of stool so it doesn't feel so full and uncomfortable – these are skid marks. OR it will just hold onto the stool, making it larger and harder all the while, so it becomes painful and scary to pass and ultimately clogs the toilet.

This cycle leads to more problems – over time, the distended rectum develops decreased sensation (sound familiar?) and ultimately the child loses the ability to feel when he has to poop. Large poop masses in distended intestines can lead to pain while eating (the kid who eats two bites of dinner, then stops because of belly pain). Poop can leak out without your child knowing because he is unaware of the sensation of the chronically-ignored-full-rectum. In addition, a full rectum will also press on the bladder changing the way the brain is sensing whether or not to void.

How can we parents prevent these problems?

Remind your child to use the bathroom upon waking up in the morning and again before leaving the house for school. Even if he says he does not have to go, tell him to “Just check- sit for 2 minutes.”

Ask your child if he uses the bathroom in school. Preschool and early elementary school teachers often have scheduled bathroom breaks, but then it's entirely up to your child to remember to use the bathroom. Encourage him to stop in the bathroom around lunch time in school.

Once home from school, remind your child to use the bathroom or “just check” if he hasn't already done so.

If you leave the house again for an after-school activity, insist that he “just check” again to decrease your chances of having to use another “foreign” toilet.

Incorporate a potty check into your child's bedtime routine, even if he went relatively recently. Ideally the healthy

bladder empties four to six times a day and the healthy bowel easily passes a stool at least once a day or every other day.

Because boys usually stand to urinate, remind them to sit at least once or twice a day, even if they don't feel the urge to poop. As a mom of three boys, trust me: they often surprise themselves.

Even middle school and high school kids can have these problems. Suggest to your older child who, for whatever reason, avoids school bathrooms, that she can stop by the school nurse's office to use the bathroom. She could also get up in the middle of class, if needed, and use the bathroom when it is most likely to be empty of other students, unlike during change-of-class time.

School cafeteria food and snacks provided at sports or other after school activities are not always healthy and can contribute to making stools too hard, so make sure to provide healthy choices at home with plenty of fruits and vegetables. Teach your kids to avoid too many cookies and other high fat foods – these are constipating foods. Encourage water-drinking. Caffeine-containing drinks (ice tea, soda) cause excess urine production and thus more stress on the bladder that your child might already be forcing to “hold it” for too long.

As for the children who frequently run to the bathroom in the daytime, but only pee a tiny bit, beware of something called urinary frequency syndrome.

Final exam

1-Does your child use the bathroom regularly, without stress or pain?

2-Can your child “go with the flow” in school as well as home, without any toilet clogging?

3-At the end of the day, is your child's underwear without urine or skid marks?

If you can answer YES to all of the above, then you and your child have passed advanced potty training!

Julie Kardos, MD and Naline Lai, MD

© 2016 Two Peds in a Pod®

Potty training 101: the nuts and bolts



A shout out to Trinity Day School in Solebury, PA where we spoke with a group of parents yesterday about the pearls and pitfalls of potty training. Today we share some of what we discussed.



At Trinity day School

“Will it ever end?” many parents ask. Time moves in slow motion for parents teaching their kids to use the potty. For those trapped in a potty training time warp, take heart. It’s been seven years since we first released our [podcast on potty training](#) and we’re proud to report that the parents who first listened to that podcast have moved onto new parenting challenges like helping with homework. For those in the midst of training, and those who are contemplating training, this post is for you.

Children master potty training typically between the ages of two and four years. Be patient, not everyone is “typical.” **More important than your child’s age is whether she shows she is developmentally ready to train.** These signs include:

- is generally agreeable/ can follow directions.
- gets a funny expression before passing urine or poop, or runs and hides, then produces a wet or soiled diaper.
- asks to be changed/ pulls on her diaper when it becomes wet or soiled- remains dry during the day time for at least two hours (look for a dry diaper after nap time.)
- NOT because grandparents are pressuring you to start

training their grandchild.

– NOT if the child is constipated—the last thing you want to do is to teach withholding to a kid who already withholds.

-NOT if a newborn sibling has just joined the family. A new baby in the house is often a time of REGRESSION, not progression. However, if your toddler begs to use the potty at this time, then by all means, allow him to try.

Make the potty a friendly place. Have a supply of books to occupy your child while she sits. Make sure her feet are secure on the floor if using a potty chair or on a stool if using the actual toilet. If using the real toilet for training, consider placing a potty training rim on the toilet seat to prevent your child from jack-knifing into the toilet. If your child is afraid of the bathroom, put the potty chair in the hall just OUTSIDE of the bathroom.

Have reasonable expectations based on age. A two year old's attention span is two minutes. Never force your child to sit on the potty. If he doesn't want to sit, then he isn't ready to train.

Your can lead a horse to water... Reward your child for sitting on the potty, even if she does not "produce." Reward by giving a high-five, verbal praise, or a small, cheap trinket such as a sticker. Do NOT promise your child a trip to Disney for potty training—otherwise, what will you do when she learns to ride a bike or tie her shoes? Plus, unless you are prepared to leave right away, the toddler/preschooler does not developmentally understand the concept of long term reward. Accept that she may simply enjoy sitting fully clothed on the potty while singing at the top of her lungs for a few weeks.

Let your child learn by imitation At home, have an open door bathroom policy so she can imitate you and her older siblings. At school, she will imitate her potty-trained classmates.

Initially, kids rarely tell their parents they “have to use the potty.” For these kids, schedule potty visits every 2-3 hours throughout the day. Do potty checks at key times such as first waking up, right before nap, and before bedtime. Be sure to spend extra time a half an hour after meals or after a warm bath. Both meals and warmth stimulate poop!

A child is potty trained when she can do the whole deal: use the potty, help wipe, help un-dress and re-dress, and wash hands.

If the child refuses to wash hands after using the potty, she is not trained. Ultimately, the goal is for her to gain independent toileting skills. However, she will need your supervision for a while.

Important note for parents of BOYS: First potty train your son to sit for ALL business. Teach him to gently press his penis downward so pee lands in the toilet and not all over the room. Once your son stands up to urinate, he may become so excited that he may never sit down again. Better to wait until he uses the potty consistently with few accidents before teaching him to stand up. Even after he begins to stand to pee, have him sit on the potty daily to allow him time to poop.

Don't be surprised if your child trains for pee before poop. In fact, many kids go through a phase when they ask for a diaper to poop in. After all, it's frightening to see/feel a chunk of your body fall into an abyss. Dump the poop from the diaper into the potty and practice waving bye-bye.

A note about night time and naps: Potty train for when your child is awake. Your child will spontaneously, without any training, stay dry at night and during naps. Some kids sleep more soundly than others and some kids are not genetically programmed to stay dry overnight until they are elementary school aged. For more information

about bed-wetting please see our post on this topic. No amount of daytime training will affect what happens during sleep. Moderate fluids right before bed and continue putting on the diapers at night until you notice that the diapers are dry when your child wakes up. After a week of dry mornings, try your child in underwear overnight. Occasional accidents are normal for years after potty training, so you might want to put a water proof liner under your child's sheets when first graduating to sleep underwear.

Disposable training pants: We like sticking to underwear while potty trainers are awake and diapers while asleep. A reluctant trainer tends to find training pants just absorbent enough that he does not care if he is wet. However, the pants are not absorbent enough to prevent rashes from stool or urine. Plus they are more expensive than underwear AND diapers. Explain to your child "sleep diapers" are perfectly acceptable until their "pee pee learns to wake them up." Use the training pants when your child is older and is mortified by the idea of a diaper or if your family is going on a long car ride and you don't want to risk urine on a car seat.

Above all: avoid power struggles. If potty training causes tears, tantrums, or confusion then STOP TRAINING, put those diapers back on, and try again a few weeks later.

After the training, keep an eye on how often he pees and poops. Older kids get "too busy" to go to the potty. Make sure he is in the habit of emptying his bladder four to six times a day and having a soft bowel movement every day or every other day.

Ultimately... you just have to go with the flow. And remember, everything eventually comes out right in the end.

Julie Kardos, MD and Naline Lai, MD

©2016, 2013 Two Peds in a Pod®

Where the boys are: raising emotionally healthy sons



photo by Lexi Logan,
www.lexilogan.com

We welcome back guest blogger Dina Ricciardo LSW, ACSW who addresses how to support the emotional health of a boy – Drs. Kardos and Lai

Your son is crying. A mad dash across the playground has led to a spectacular trip and fall, complete with a bloody knee and hands full of dirt. Part of you wants to hold him on your lap and console him until he stops crying. The other part of you wants to firmly wipe away his tears and tell him to be brave. Which part of you is right?

In a world where there is a great deal of emphasis placed on the emotional health of girls, our boys are frequently overlooked. While girls are typically encouraged to develop and express a broad range of emotions, boys are socialized from a young age to suppress their feelings. As a result, many boys and men struggle to express fear or sadness and are unable to ask for help. It is time for us adults to stop perpetuating stereotypes and myths about manhood, and help each other raise emotionally healthy boys. Here are five ways for us to do so:

Make his living environment a safe space to express emotions.

Give your son permission to express *all* of his feelings. Boys typically do not have the freedom to show the full range of their emotions in school and out in the world, so it is essential that they have that freedom at home. Nothing should be off limits, as long as feelings are expressed in a manner that is not destructive.

Expose him to positive male role models. Boys need to be exposed to positive male figures who can indoctrinate them into their culture and teach them how to be men. It is an important rite of passage in a boy's development. Take a look around your social ecosystem and ask yourself, "Who would be good for my son?" Other parents, coaches, teachers, and pastors are examples of individuals who can play a positive role in his life.

Understand your unique role. Each parent plays a unique role in the development of a son, and that role changes over time. A mother is a son's first teacher about love and what it looks like, and this dynamic can breed a particular kind of closeness. As a boy grows and begins to develop his sexuality, however, it is natural for him to pull away a bit from his mother and turn more towards his father for guidance. While this distance can be unsettling for mom, it marks a new phase in a son's relationship with his father, who typically provides a sense of security and authority in a family as well

as support for a boy's developing identity. Mothers still play an important role, but that role may look different. As parents, it is important to re-evaluate what our sons need from us at each stage of their development.

Look at the world with a critical eye. Our culture not only glorifies violence, it equates vulnerability in males with weakness and attempts to crush it. That does not mean we have to accept this paradigm. Talk honestly with your son about how and when to be gentle and compassionate, educate him on how the world view softness in men, and never tolerate anyone shaming him when he exhibits these traits. There is no shame in showing vulnerability, it is actually an act of courage.

Take a look in the mirror. Whether you are a mother or a father (or both), be honest with yourself: what are your beliefs about manhood? Do you feel safe expressing all of your feelings, or are some of them off-limits? If you are perpetuating negative stereotypes about men or are not comfortable with a full range of emotions, then your son will follow in your footsteps. Regardless of our own gender, we cannot expect our children to be comfortable with their feelings if we are not comfortable with our own.

There are times when insuring the emotional health of your son will feel like an uphill battle. Keep the conversation open, and do not be afraid to talk with others about the dilemmas of boyhood and manhood. And if you are looking for an answer to the playground dilemma, then I will tell you that both parts of you are right. Sometimes our sons need loving compassion, and sometimes they need a firm nudge over the hump. You know your child better than anyone else, so it is up to you to decide which approach to use and when.

Dina Ricciardi, LSW, ACSW

©2016 Two Peds in a Pod®

Dina Ricciardi is a psychotherapist in [private practice](#)

treating children, adolescents, and adults in Doylestown, PA. She specializes in disordered eating and pediatric and adult anxiety, and is also trained in Sandtray Therapy. Ricciardi is a Licensed Social Worker and a member of the Academy of Certified Social Workers. She can be reached at dina@nourishcounseling.com.

When your child says, “My belly hurts”



This week Two Peds joined Kelley on her blog Happy Health Kids as she talked about the dreaded phrase, “My belly hurts.”

If I crunched the numbers on how often my kids have uttered certain phrases, “my belly hurts,” ranks pretty darn high. So common is this refrain, and typically uneventful the outcome, that there’s a cry-wolf quality to it; I typically point my child towards the pantry or bathroom and go about my day. But

sometimes, a stomachache persists, and then figuring out the cause can be like falling into a rabbit hole (and equally unpleasant)...[click here to continue](#)