

# Stealthy Salmonella! Not just in your eggs



Raw chicken left out overnight—Dr. Lai's recipe for Salmonella

These days it seems that the bacteria *Salmonella* is lurking everywhere. Last month's egg recall for possible *Salmonella* contamination affected over 200 million eggs, but *Salmonella* is not just in eggs. In the last few months, dried coconuts and even guinea pigs (as pets, not as food!) have caused people gastrointestinal misery.

Nontyphoidal *Salmonella* usually causes fever and crampy diarrhea. Sometimes the stools contain blood. This stomach bug mainly lurks in raw poultry, raw eggs, raw beef, and unpasturized dairy products. Luckily, *Salmonella* does not jump up and attack humans. In general, people are safe from disease as long as they do not eat salmonella-infested food. But children below the age of five often put their hands in their mouths and can acquire *Salmonella* after touching a

contaminated source.

Reptiles such as lizards and turtles can carry *Salmonella* in their stool and are not recommended as pets for young children. Turtles that are four inches or smaller (about the size of a deck of playing cards) are most likely to harbor the bacteria. As a preschooler, Dr. Kardos remembers that her tiny pet turtle suddenly disappeared. Her parents told her that "Her pet would be happier if it went outside to the stream to swim with the other turtles." In retrospect, Dr. Kardos thinks her pediatrician dad was worried about *Salmonella* and made the turtle magically disappear.

Even cute little chicks can be problematic. *Salmonella* carried in the gut of a chick can contaminate the entire surface of a chick. So, although kissing and cuddling a chick makes for a good Instagram post, discourage your children from doing so.

Unfortunately, you cannot depend on a warning stench arising from your lunch to warn you that it is inedible. *Salmonella* often hides in food and it is difficult to tell what is or is not contaminated. So how can you prevent your kids from catching *Salmonella*?

Luckily *Salmonella* is killed by heat and bleach. Even if an otherwise fine-looking and odorless raw egg has *Salmonella*, adequate cooking will destroy the bacteria. Gone are the days when parents can feed kids soft boiled eggs in a silver cup, have kids wipe up with toast the yolk from a sunny-side up egg, add a raw egg to a milkshake, or let their kids lick the left-over cake batter from the mixing bowl. Instead, cook hardboiled eggs until the yolks are green and crumble, and make sure your scrambled eggs aren't runny. Wash all utensils well. The disinfecting solution used in childcare centers of  $\frac{1}{4}$  cup bleach to 1 gallon water works well to sanitize counters. Do not keep perishable food, even if it is cooked, out at room temperature for more than two hours. And wash, wash, wash your hands.

A mom once called us frantic because her child had just happily eaten a half-cooked chicken nugget. What if this happens to your child? Don't panic. Watch for symptoms – the onset of diarrhea from *Salmonella* is usually between 12 to 36 hours after exposure but can occur up to three days later. The diarrhea can last up to 5-7 days. If symptoms occur, the general recommendation is to ride it out. Prevent dehydration by giving plenty of fluids. My simple rule to prevent dehydration is that more must go in than comes out.

Although of unproven benefit, antibiotic treatment may be considered if your child is at risk for overwhelming infection, including infants younger than three months old and those with abnormal immune systems (cancer, HIV, Sickle Cell disease, kids taking daily steroids for other illnesses) or those with chronic gastrointestinal tract diseases\*. Using antibiotics to treat a typical case of salmonella not only promotes general antibiotic resistance, but also does not shorten the time frame for the illness. In fact, the medication can prolong how long your child carries the germ in his stool.

Pictured above is a pot of chicken Dr. Lai accidentally left out overnight one warm summer night. Yuck.


Naline Lai, MD and Julie Kardos, MD

©2018 Two Peds in a Pod®

\**Red Book*, 2015 Report of the Committee on Infectious Diseases, American Academy of Pediatrics

---

# How to treat your kid's allergies: sorting out over the counter medications



Gepetto always said his son had allergies, but the villagers knew better

It's not your imagination. This is a particularly bad spring allergy season. We didn't need media outlets to tell us that there are more itchy, sneezy, swollen eyed kids out there this year.

It is worth treating your child's allergy symptoms- less itching leads to improved sleep, better ability to pay attention in school, improved overall mood, and can prevent asthma symptoms in kids who have asthma in addition to their nose and eye allergies.

Luckily, nearly every allergy medication that we wrote prescriptions for a decade ago is now available over-the-counter. As you and your child peer around the pharmacy through itchy blurry eyes, the displays for allergy medications for kids can be overwhelming. Should you chose the medication whose ads feature a bubbly seven-year-old girl kicking a soccer ball in a field of grass, or the medication whose ads feature a bubbly ten-year-old boy roller blading? Its it better to buy a “fast” acting medication or medication that promises your child “relief?”

Here is a guide to sorting out your medication choices:

**Oral antihistamines:** Oral antihistamines differ mostly by how long they last, how well they help itchiness, and their side effect profile. During an allergic reaction, antihistamines block one of the agents responsible for producing swelling and secretions in your child’s body, called histamine. Prescription antihistamines are not necessarily “stronger.” In fact, at this point there are very few prescription antihistamines. The “best” choice is the one that alleviates your child’s symptoms the best. As a good first choice, if another family member has had success with one antihistamine, then genetics suggest that your child may respond as well to the same medicine. Be sure to check the label for age range and proper dosing.

**First generation antihistamines work well at drying up nasal secretions and stopping itchiness** but don’t tend to last as long and often make kids very sleepy. **Diphenhydramine (brand name Benadryl)** is the best known medicine in this category. It lasts only about six hours and can make people so tired that it is the main ingredient for many over-the-counter adult sleep aids. Occasionally, kids become “hyper” and are unable to sleep after taking this medicine. Opinion from Dr. Lai: dye-free formulations of diphenhydramine are poor tasting. Other first generation antihistamines include **Brompheniramine (eg. brand names Bromfed and Dimetapp)** and **Clemastine (eg.**

brand name Tavist).

**Second and third generation antihistamines cause less sedation and are conveniently dosed only once a day. Cetirizine (eg. brand Zyrtec)** causes less sleepiness and it helps itching fairly well. Give the dose to your child at bedtime to further decrease the chance of sleepiness during the day. **Loratadine (brand name Alavert, Claritin)** causes less sleepiness than cetirizine. **Fexofenadine (brand name Allegra)** causes the least amount of sedation. The liquid formulations in this category tend to be rather sticky, the chewables and dissolvables are favorites among kids. For older children, the pills are a reasonable size for easy swallowing.

**Allergy eye drops:** Your choices for over-the-counter antihistamine drops include **ketotifen fumarate (eg. Zatidor and Alaway)**. For eyes, drops tend to work better than oral medication. Avoid products that contain vasoconstrictors (look on the label or ask the pharmacist) because these can cause rebound redness after 2-3 days and do not treat the actual cause of the allergy symptoms. Contact lenses can be worn with some allergy eye drops- check the package insert, and avoid wearing contacts when the eyes look red. Artificial tears can help soothe dry itchy eyes as well.

**Allergy nose sprays:** Simple nasal saline helps flush out allergens and relieves nasal congestion from allergies. **Flonase**, which used to be available by prescription only, is a steroid allergy nose spray that is quite effective at eliminating symptoms. It takes about a week until your child will notice the benefits of this medicine. Even though this medicine is over-the-counter, check with your child's pediatrician if you find that your child needs to continue with this spray for more than one allergy season of the year. Day in and day out use can lead to thinning of the nasal septum. Avoid the use of nasal decongestants (e.g., Afrin, Neo-Synephrine) for more than 2-3 days because a rebound runny nose called rhinitis medicamentosa may occur.

**Oral Decongestants** such as phenylephrine or pseudoephedrine can help decrease nasal stuffiness. This is the “D” in “Claritin D” or “Allegra D.” However, their use is not recommended in children under age 6 years because of potential side effects such as rapid heart rate, increased blood pressure, and sleep disturbances.

Some of the above mentioned medicines can be taken together and some cannot. Read labels carefully for the active ingredient. Do not give more than one oral antihistamine at a time. In contrast, most antihistamine eye drops and nose sprays can be given together along with an oral antihistamine.

If you are still lost, call your child’s pediatrician to tailor an allergy plan specific to her needs.

**The best medication for kids?** Get the irritating pollen off your child. Have your allergic child wash her hands and face as soon as she comes in from playing outside so she does not rub pollen into her eyes and nose. know that spring and summer allergens/pollen counts are highest in the evening, vs fall allergies where counts are highest in the mornings. Rinse outdoor particles off your child’s body with nightly showers. Filter the air when driving in the car and at home: run the air conditioner and close the windows to prevent the “great” outdoors from entering your child’s nose. If you are wondering about current pollen counts in your area, scroll down to the bottom of many of the weather apps to find pollen counts or log into the American Academy of Allergy Asthma and Immunology’s website.

Naline Lai MD and Julie Kardos, MD

© 2018 Two Peds in a Pod®

---

# Contribute to our Two Peds Mother's Day post!



Dr. Kardos, on a visit home from medical school, with her mom and grandmothers, 1991.

A flash of surprise spread across her face. "You mean my mother was right? I can't believe it!" the mom in our office exclaimed.

Many times as we dispense pediatric advice, the parent in our office realizes that their own mother had already offered the same suggestions.

This Mother's Day, we're asking readers for anecdotes about times where maybe, just maybe, your mom or your grandmother was right after all. If you have a photo available of your mom or grandmother with your child that you don't mind sharing as well, we would love to post them along with your anecdotes this Mother's Day.

Please send them along to us at [twopedsinapod@gmail.com](mailto:twopedsinapod@gmail.com) before Mother's Day weekend.

Naline Lai, MD and Julie Kardos, MD



# Kids are vaping: e-cigarettes



Photo by American Academy of Pediatrics, Parent Plus 2018

It's time for another Two Peds in a Pod photo quiz.

The question: What's depicted in this photo?

If you answered: a pen, a thumb nail drive, or an asthma inhaler, you would be wrong.

Kids use these devices, which purposely look like common innocuous objects, to inhale electronic cigarettes (e-cigarettes). Vaping, also called “Juuling” and an even more concentrated form of vaping, called “dripping,” is unfortunately popular among teens. It’s unhealthy: the stuff that the kids are inhaling contains nicotine and other chemicals.

Ask your middle schooler or high schooler. They most likely have seen these devices if they have not actually used one.

Parents need to know kids are vaping in school as well as outside of school. Unlike conventional cigarettes, it’s easy for the kids to hide: no smoky smell, no cigarette cartons. The vaping liquid or “e-juice” comes in all kinds of “kid friendly” flavors such as gummy bear, fruit, or chocolate, and the devices needed to inhale them look like items in every kid’s pencil case or backpack.

It’s easy for kids to get the e-juice on the internet because online stores don’t always ask for proof of age (legal age to buy is 18 years in the US). Most e-juices contain nicotine, which is addictive. Emerging data show that kids who vape are more likely to go on to use regular cigarettes than kids who do not vape.

Bottom line: Vaping, or using electronic cigarettes, is unhealthy and addictive, and startlingly easy for kids to hide.

Julie Kardos, MD and Naline Lai, MD

©2018 Two Peds in a Pod®

---

# Worry wart: how to treat a wart



Nope, warthogs don't actually have warts. But kids often do!


Emma's dad and I both peered at the filamentous growth dangling from his nine year old's right nostril. "Yes," I said, "it's definitely a wart."

Emma's dad offered, "When I was a kid, I heard the way to get rid of a wart was to cut a potato in half, rub it on the wart, and bury the potato in the backyard. Legend had it, by the time the potato disintegrates, the wart will be gone."


"I wish it were so easy," I replied.

Warts are caused by skin-dwelling viruses. On the feet, warts can sometimes be mistaken for calluses. One distinguishing feature is that warts sit in the skin like this:





Fine “feeder” blood vessels extend from the wart into the skin. Therefore, if you scrape off the top layer of a wart, a dotted pattern usually appears from above. The dots will not appear in a callus. View from above:



There are simply no glamorous ways to get rid of warts. Most treatment modalities destroy warts by pulverizing the home they live in, a.k.a. your skin. Your doctor may be armed with various agents such as liquid nitrogen or dimethyl ether propane, which produces a chemical “freeze” and dries up the wart. Another agent called cantharidin (otherwise known as “beetle juice”) is a caustic liquid derived from the blister beetle. Application of beetle juice causes the warts to blister.

Some doctors will even manually take a scalpel and cut out the

warts.

Like I said, there are no glamorous treatments. However, more gentle creams which stimulate the immune system, such as Imiquimod (Aldara) show some promise in children. Other compounds such as 5-fluorouracil can be topically applied or injected and treatments such as pulsed dye laser therapy have mixed reviews.

Over-the-counter remedies exist in a milder form. Commonly used wart removers such as Compound W, Dr Scholl's Clear Away Wart, and Duofilm all contain salicylic acid. The acid slowly dries up the warts. When applying salicylic acid, after a few applications make sure you peel the dead crusty top layer off the wart. Without peeling, future medicine will not reach the wart. These methods can take weeks to months to work, but they do work.

And don't forget the duct tape. Duct tape, the great all-purpose household item, has also been shown to speed up the resolution of warts. Scientists hypothesize the constant presence of the adhesive somehow stimulates a natural immune response. If you try duct tape, have your child wear the duct tape over the wart for several days in a row and then give a day off. If the wart is on a hand or foot, the tape tends to fall off during the day: just re-apply some tape at bedtime. Effects should be seen within a couple of months if not sooner. Now, the original study that showed duct tape was helpful, was followed by a study which showed duct tape was not helpful. Some hypothesize that the results differ because silver sticky duct tape was used in the initial study, while the later study used less sticky duct tape. So be sure to use the old-fashioned silver duct tape.

The prevention of warts is tricky. Some people just seem genetically predisposed. However, your best bet for keeping warts away is to keep your child's skin as healthy as possible. Warts tend to gravitate towards areas of skin

broken down by friction such as feet or fingers. Liberally apply moisturizing creams daily to prone areas. After a summer of wearing flip-flops and walking on the rough cement by the side of a swimming pool in bare feet, many children end up with warts on the bottom of their feet. I know a teen whose warts on the tips of her fingers stemmed from months of guitar strumming.

Turns out that even without treatment, 60% percent or more of all warts will disappear spontaneously within two years.

Coincidentally, I think that's also the time it takes for a potato half to disintegrate.

Naline Lai , MD and Julie Kardos, MD

© 2009, 2018 TwoPeds in a Pod®

---

## **“Ya Gotta Have Heart!” Heart Murmurs Explained**



## Conversation hearts murmuring

When the Tin Man was a child in Oz, I'm sure his pediatrician never told his parents, "Has anyone ever said your child has a heart murmur? I hear one today."

I know that when I tell parents about a heart murmur in their child, their hearts skip and jump. But not all heart murmurs are bad.

## What is a heart murmur?

A heart murmur is an extra sound that we pediatricians hear when we listen to a child's heart with a stethoscope. A normal heart beat sounds like this: "lub, dub. lub, dub. lub, dub." A heart murmur adds a whooshing sound. So what we hear instead is "lub, *whoosh*, dub" or "lub, dub, *whoosh*."

The “whoosh” is usually caused by blood flowing through a relatively narrow opening somewhere in or around the heart. Think of your blood vessels and heart like a garden hose. If you run the water (blood) very hard, or put a kink or cut a hole in the hose, the whoosh of the water grows louder in those locations.

## **Heart murmurs signal different issues at different ages.**

In a newborn, some types of heart murmurs are expected. Normal newborn hearts contain extra holes that close up after the first hours or days of birth. One type of murmur occurs as the infant draws in his first breath and holes in the heart, present inside the womb, begin to seal. As the holes get narrower, we sometimes hear the “whoosh” of blood as it flows through the narrowing opening. Then these holes close completely and the murmur goes away.

However, some murmurs in infancy signal “extra holes” in the heart. As pediatricians, we experience our own heart palpitations when moms want to leave the hospital early with their infants who are less than 48 hours old. We worry because many infants who have abnormal hearts may not develop their abnormal heart murmurs and other signs of heart failure until the day or two after birth.

Preschool and early school-age children often develop “innocent” heart murmurs. “Innocent” implies that extra blood flows through their hearts, but the hearts are structurally normal. These murmurs are fairly common and can run in families. However, there are some significant heart problems which do not surface until this age. For this reason, remember to schedule those yearly well child checkups.

For teens, during the pre-participation sports physical, pediatricians listen carefully for a murmur that may indicate that an over grown heart muscle has developed.



## **What else can cause a heart murmur?**

Holes are not the only culprit behind a murmur. The whoosh sound can also arise when a person is anemic and blood flows faster than normal. In anemic kids, the blood flows faster because it lacks enough oxygen-carrying red blood cells and the heart needs to move blood faster in order to supply oxygen to the body. The most common cause for anemia is a lack of eating enough iron-containing foods. Subsequently, we hear these flow murmurs in children whose diets lack iron, in teenagers who grow rapidly and quickly use up their iron stores, and in girls who bleed too much at each period. Replenishing the iron level makes a heart murmur from anemia go away.

Even a simple fever can cause a heart murmur on physical exam. The murmur goes away when the fever goes away.

Pediatric health care providers can often distinguish between “innocent” heart murmurs and not-so- innocent heart murmurs by the sound of the murmur itself (not all “whooshes” sound alike). If any question exists, your child will be referred for more testing, which could include a chest x-ray, an EKG (electrocardiogram), an ECHO (echocardiogram, or ultrasound of the heart), or evaluation by a pediatric cardiologist.

## **If your child’s pediatrician tells you that your child has a heart murmur, “take heart.”**

Many times a murmur comes and goes or just becomes part of your child’s baseline physical exam. Even if your child has a serious heart problem, most cases respond well to medication, surgery, or both. While not all heart problems cause heart murmurs, and while not all murmurs signal heart problems, the presence of a heart murmur in a child can signal that your child needs further testing.

Unless, of course, your child is the Tin Man. In this case, extra sounds indicate that your child needs more oil!

Julie Kardos, MD and Naline Lai, MD

©2010, 2018 Two Peds in a Pod®

---

# Raising emotionally healthy boys



photo by Lexi Logan,  
[www.lexilogan.com](http://www.lexilogan.com)

*The recent Parkland shooting in Florida is causing many to wonder how to support the emotional health of boys in their families and communities. We welcome therapist Dina Ricciardo's words of wisdom— Drs. Kardos and Lai*

Your son is crying. A mad dash across the playground has led to a spectacular trip and fall, complete with a bloody knee and hands full of dirt. Part of you wants to hold him on your lap and console him until he stops crying. The other part of you wants to firmly wipe away his tears and tell him to be brave. Which part of you is right?

In a world where there is a great deal of emphasis placed on the emotional health of girls, our boys are frequently overlooked. While girls are typically encouraged to develop and express a broad range of emotions, boys are socialized from a young age to suppress their feelings. As a result, many boys and men struggle to express fear or sadness and are unable to ask for help. It is time for us adults to stop perpetuating stereotypes and myths about manhood, and help each other raise emotionally healthy boys. Here are five ways for us to do so:

**Make his living environment a safe space to express emotions.**

Give your son permission to express *all* of his feelings. Boys typically do not have the freedom to show the full range of their emotions in school and out in the world, so it is essential that they have that freedom at home. Nothing should be off limits, as long as feelings are expressed in a manner that is not destructive.

**Expose him to positive male role models.** Boys need to be exposed to positive male figures who can indoctrinate them into their culture and teach them how to be men. It is an important rite of passage in a boy's development. Take a look around your social ecosystem and ask yourself, "Who would be good for my son?" Other parents, coaches, teachers, and pastors are examples of individuals who can play a positive role in his life.

**Understand your unique role.** Each parent plays a unique role in the development of a son, and that role changes over time. A mother is a son's first teacher about love and what it looks

like, and this dynamic can breed a particular kind of closeness. As a boy grows and begins to develop his sexuality, however, it is natural for him to pull away a bit from his mother and turn more towards his father for guidance. While this distance can be unsettling for mom, it marks a new phase in a son's relationship with his father, who typically provides a sense of security and authority in a family as well as support for a boy's developing identity. Mothers still play an important role, but that role may look different. As parents, it is important to re-evaluate what our sons need from us at each stage of their development.

**Look at the world with a critical eye.** Our culture not only glorifies violence, it equates vulnerability in males with weakness and attempts to crush it. That does not mean we have to accept this paradigm. Talk honestly with your son about how and when to be gentle and compassionate, educate him on how the world view softness in men, and never tolerate anyone shaming him when he exhibits these traits. There is no shame in showing vulnerability, it is actually an act of courage.

**Take a look in the mirror.** Whether you are a mother or a father (or both), be honest with yourself: what are your beliefs about manhood? Do you feel safe expressing all of your feelings, or are some of them off-limits? If you are perpetuating negative stereotypes about men or are not comfortable with a full range of emotions, then your son will follow in your footsteps. Regardless of our own gender, we cannot expect our children to be comfortable with their feelings if we are not comfortable with our own.

There are times when insuring the emotional health of your son will feel like an uphill battle. Keep the conversation open, and do not be afraid to talk with others about the dilemmas of boyhood and manhood. And if you are looking for an answer to the playground dilemma, then I will tell you that both parts of you are right. Sometimes our sons need loving compassion, and sometimes they need a firm nudge over the hump. You know

your child better than anyone else, so it is up to you to decide which approach to use and when.

Dina Ricciardi, LSW, ACSW

©2016, 2018 Two Peds in a Pod®

*Dina Ricciardi is a psychotherapist in [private practice](#) treating children, adolescents, and adults in Doylestown, PA. She specializes in disordered eating and pediatric and adult anxiety, and is also trained in Sandtray Therapy. Ricciardi is a Licensed Social Worker and a member of the Academy of Certified Social Workers. She can be reached at [dina@nourishcounseling.com](mailto:dina@nourishcounseling.com).*

---

## **When a pet dies**



Photo by Lexi Logan

*We welcome Bereavement Counselor Amy Keiper-Shaw who shares with us how to discuss the death of a pet with your child.*

*—Drs. Lai and Kardos*

When I first graduated from college I worked as a nanny. One day the mom shared with me that their family goldfish recently died. As this was her daughter's first experience with death, we schemed for nearly 20 minutes to find the best way to talk to her child. The mom and I thought it could be an excellent teaching moment.

We pulled the girl away from her playing to explain that the fish had died. We told the girl we'd help her have a funeral if she wanted, and we would find a box (casket) to bury the fish so she could say her goodbyes. We explained what a casket

was and what a funeral was in minute detail. After our monologue we stopped, we asked if she had any questions.

After a slight pause she asked, "Can't we just flush it?"

The lesson I learned from that experience, and still use to this day, is to keep things simple, and know my audience. Sometimes as parents we overcompensate for our own fears and make situations more challenging than they need to be.

**Here are some tips on how to talk to your children about pet loss:**

**Tell your child about the death, and then pause.** Ask her what she thinks death means before moving on with further explanations. This will help you know if she has questions or if she has enough information for the moment. Children often need a small amount of information initially and will later come back to you several times later to ask more questions after they process the information.

**Remember to express your own grief,** and reassure your child that many different feelings are ok. Be sure to allow children to express their feelings. If your child is too young to express herself verbally, give her crayons and paper or modeling clay too help express grief.

**Avoid using clichés such as: Fluffy "went to sleep."** Children may develop fears of going to bed and waking up. The phrase "God has taken" the pet could create conflicts in a child and she may become angry at a higher power for making the pet sick, die, or for "taking" the pet from them.

**Be honest.** Hiding a death from a child can cause increased anxiety. Children are intuitive and can sense if something is wrong. When the death isn't explained they make up their own explanation of the truth, and this is often much worse than the reality of what occurred.

Children are capable of understanding that life must end for

all living things. Support their grief by acknowledging their pain. The death of a pet can be an opportunity for a child to learn that adult caretakers can be relied upon to extend comfort and reassurance through honest communication.

## **Developmental Understanding of Death**

### **Two and three-year-olds**

Often consider death as sleeping, therefore tell them the pet has **died** and will not return.

Reassure children that the pet's failure to return is unrelated to anything the child may have said or done (magical thinking).

A child at this age will readily accept another pet in the place of a loved one that died.

### **Four, five, and six-year-olds**

These children have some understanding of death but also a hope for continued living (a pet may continue to eat, play & breathe although deceased).

They can feel that any anger that they had towards the pet may make them responsible for the pet's death ("I hated feeding him everyday").

Some children may fear that death is contagious and could begin to fear their own death or worry about the safety of their parents.

Parents may see temporary changes in their child's bladder/bowels, eating, and sleeping.

Several brief discussions about the death are more productive than one or two prolonged discussions.

### **Seven, eight, and nine-year-olds**

These children have an understanding that death is real and irreversible.



Although, to a lesser degree than a four, five, or six-year-old, these children may still possibly fear their own death or the death of their parents.

May ask about death and its implications (Will we be able to get another pet?).

Expressions of grief may include: somatic concerns, learning challenges, aggression, and antisocial behavior. Expression may take place weeks or months after the loss.

### **Adolescents**

Reactions are similar to an adult's reaction.

May experience denial which can take the form of lack of emotional display so they could be experiencing the grief without outwards manifestations.

### **Resources:**

Petloss.com— a gentle and compassionate website for pet lovers who are grieving the death or an illness of a pet- they have a Pet Loss Candle Ceremony every week

Your local veterinarian- often your veterinarian has or knows of a local pet loss group

Handsholdinghearts.org— our group of counselors offer grief support to children, teens, and their families centered in Bucks County Pennsylvania.

### **Books on pet loss for children:**

*Badger's Parting Gifts* (children) by Susan Varley

*Lifetimes* by Brian Mellonie & Robert Ingpen

*The Tenth Good Thing About Barney* (children) by Judith Viorst

Amy Keiper-Shaw, LCSW, QCSW, GC-C

©2013, 2018 Two Peds in a Pod®

Amy Keiper-Shaw is a licensed grief counselor who holds a Masters Degree in clinical social work from the University of

Pennsylvania. For over a decade she has served as a bereavement counselor to a hospice program and facilitates a bereavement camp for children. She directs Handsholdinghearts, a resource for children who have experienced a significant death in their lives.

---

## Home remedies for dry, chapped hands



Raw hands- recognize your kid?

Even when it isn't flu season, we pediatricians wash our hands about sixty times a day, maybe more. This frequent washing, in combination with cold winter air, leads to dry, chapped hands. Here are the hands of a patient. Do your children's hands look like these?

**To prevent dry, chapped hands:**

- **Don't stop washing your hands**, but do use a moisturizer afterwards. Also use warm but not hot water. Hot water removes protective oils from skin.
- According to the American Academy of Dermatology, **hand sanitizer can prevent the drying** that accompanies frequent hand washing. However, we can tell you from experience that once your hands are already chapped and cracked, the alcohol content in the sanitizers stings sensitive skin. So if your child's hands are already chapped, stick with water and soap.
- **Wear gloves or mittens** as much as possible outside even if the temperature is above freezing. Remember chemistry class—cold air holds less moisture than warm air and therefore is unkind to skin. Gloves will prevent some moisture loss. Having difficulty convincing your child to wear gloves? Point out that refrigerators are kept around 40 degrees Fahrenheit or below. Tell your kids that if they wouldn't sit inside a refrigerator without layers, then it would be wise to wear gloves.
- Before exposure to any possible irritants such as the chlorine in a swimming pool, **protect the hands by layering heavy lotion (e.g. Eucerin cream) or petroleum based product (e.g. Vaseline or Aquaphor) over the skin.**

#### **To rescue dry, chapped hands:**

- Prior to bedtime, smother hands in **1% hydrocortisone ointment**. Avoid the cream formulation. Creams tend to sting if there are any open cracks. Take old socks, cut out thumb holes and have your child sleep at night with the sock on his hands. Repeat nightly for up to a week. Alternatively, for mildly chapped hands, use a **petroleum oil based product such as Vaseline or Aquaphor** in place of the hydrocortisone.
- If your child has underlying eczema, **prevent your child from scratching his hands**. An antihistamine taken orally such as diphenhydramine (Benadryl) or cetirizine (Zyrtec) will take the edge off the itch. Keep his nails trimmed to avoid further damage from scratching.

- **For extremely raw hands**, your child's doctor may prescribe a stronger cream and if there are signs of a bacterial skin infection, your child's doctor may prescribe an antibiotic.

Happy moisturizing. Remember smearing glue on your hands and then peeling off the dried glue? It's not so fun when your skin really is peeling.

Naline Lai, MD and Julie Kardos, MD

© 2009, updated 2019, Two Peds in a Pod®

---

## How to explain scary news to children



In light of the recent school shooting in Parkland, Florida, you may be left wondering if, and how, to explain this or other tragedies to your children.

Understand that kids sense your emotions even if you don't tell them. Not telling them about an event may make them

concerned that they are the cause for your worried hushed conversations. Break away from your discussion with adults to say, “ Do you know what we are talking about? We are not talking about you.”

Even though an event may be far away, media makes it seem as if it happened next door, and sooner or later your children will see or hear about it. Tell the facts in a straight forward, age appropriate manner. Answer questions and don't be afraid to answer with an “I don't know.” Preschoolers are concrete in their thinking—dragons are real and live under their bed, so don't put any there that do not exist. For a preschooler a simple “Mom is sad because a lot of people got hurt,” will suffice. Young school age kids will want to know more details. And be prepared to grapple with more high level questions from teens.

Look for the helpers. Mr. Rogers who hosted Mister Roger's Neighborhood for 30 years, tells this story about seeing scary things on the news: “My mother would say to me, ‘Look for the helpers. You will always find people who are helping.’ To this day, especially in times of ‘disaster,’ I remember my mother's words, and I am always comforted by realizing that there are still so many helpers-so many caring people in this world.”

What if the kids ask, “Will that happen here?” or “Why did that happen?” Again, reassure in a simple straight forward manner. For instance you can say, “Many people are working hard to prevent something like that here.” Consider answering the question with a question. Asking “What do you think?” will give you an idea of exactly what your child fears. You can also reach out to other family supports for help with answers. Say to your child, “I wonder what our minister or school counselor has to say about this, let's ask.”

Routine is reassuring to children, so turn off the background 24 hour television and internet coverage and make dinner, take them to sports activities, and get the homework done.

Give your kids something tangible to do to be helpful. Help them set up a coin donation jar at school or put aside part of their allowance for a donation.

If your child seems overly anxious and fearful, and her worries are interfering with her ability to conduct her daily activities, such as performing at school, sleeping, eating, and maintaining strong relationships with family and friends, then seek professional help.

For more advice on this topic, please see this American Academy of Pediatrics recommendation for parents. Also, the following is a firearms safety message for parents, from the AAP:

- Young children are curious. They are often unable to remember or follow safety rules. Teens are impulsive, and naturally tend to be moody. When combined with access to firearms, the consequences can be **tragic and permanent**.
- Many homes have guns – which is why **you have to ask about guns when your child visits another home**. Depending on location, 18 percent to 64 percent of U.S. homes have firearms.
- **If there is a gun in the home, there's a good chance a child living there will know where it is**. In a recent study, 39 percent of parents erroneously believed their children did not know where their gun is stored, and 22 percent wrongly believed their child never handled a gun.
- **The safest home for children and teens is one without guns**. If you do have a gun, you can greatly reduce the risk of a child being injured or killed by storing the gun unloaded and locked, with the ammunition locked in a separate place.

Parent your children so they feel secure in themselves and secure in the world around them. You may not hold the answers to why a tragedy strikes, but you do hold the ability to

comfort and reassure your children.

Naline Lai, MD and Julie Kardos, MD

© 2018 Two Peds in a Pod®, adapted from our 2015 post