

Tick removal and prevention



The classic bullseye rash of Lyme

Kids are running about and so are the ticks. We're talking about tick removal and prevention of tick bites in the Children's Hospital of Philadelphia Tip of the Week!

Julie Kardos, MD and Naline Lai, MD

©2020 Two Peds in a Pod®

Lyme Disease...it's back



The classic bullseye rash of Lyme

Just like last year, experts are predicting more Lyme disease. While it used to be a pesky disease only in our midatlantic/Lyme Connecticut area of the world, Lyme continues to appear across the northeast and has been reported on the west coast of the United States. According to the American Academy of Pediatric's Redbook, about fifty percent of reported Lyme disease is during June and July. We've already had children come to our office with tick bites concerns, so here's an update:

Lyme disease is spread to people by blacklegged ticks. Take heart- even in areas where a high percentage of blacklegged

ticks carry the bacteria that causes Lyme disease, the risk of getting Lyme from any one infected tick is low. Most of the little critters **DON'T** carry Lyme disease... but there are an awful lot of ticks out there. Blacklegged ticks are tiny and easy to miss on ourselves and our kids. In the spring, the ticks are in a baby stage (nymph) and can be as small as a poppy seed or sesame seed. To spread disease, the tick has to be attached and feeding on human blood for more than 36 hours, and engorged.

In areas in the United States where Lyme disease is prevalent (New England and Mid-Atlantic states, upper Midwest states such as Minnesota and Wisconsin, and California), parents should be vigilant about searching their children's bodies daily for ticks and for the rash of early Lyme disease. Tick bites, and therefore the rash as well, especially like to show up on the head, in belt lines, groins, and armpits, but can occur anywhere. When my kids were young, I showered them daily in summer time not just to wash off pool water, sunscreen, and dirt, but also for the opportunity to check them for ticks and rashes. Now that they are older I call through the bathroom door periodically when they shower: "Remember to check for ticks!" Read our post on [how to remove ticks](#) from your kids.

"I thought that Lyme is spread by deer ticks and deer are all over my yard." Nope, it's not just Bambi that the ticks love. Actually, there are two main types of blacklegged ticks, *Ixodes Scapularis* and *Ixodes Pacificus*, which both carry Lyme and feed not only on deer, but on small animals such as mice. (Fun fact: *Ixodes Scapularis* is known as a deer tick or a bear tick.)

Most kids get the classic rash of Lyme disease at the site of a tick bite. The rash most commonly occurs by 1-2 weeks after the tick bite and is round, flat, and red or pink. It can have some central clearing. The rash typically does not itch or hurt. **The key is that the rash expands to more than 5 cm**, and can become quite large as seen in the above photo. This

finding is helpful because if you think you are seeing a rash of Lyme disease on your child, you can safely wait a few days before bringing your child to the pediatrician because the rash will continue to grow. The Lyme disease rash does not come and then fade in the same day, and the small (a few millimeters) red bump that forms at the tick site within a day of removing a tick is not the Lyme disease rash. Knowing that a rash has been enlarging over a few days helps us diagnose the disease. Some kids have fever, headache, or muscle aches at the same time that the rash appears.

If your child has early localized Lyme disease (just the enlarging red round rash), the diagnosis is made by having a doctor examine your child. Your child does not need blood work because it takes several weeks for a person's body to make antibodies to the disease, and blood work checks for antibodies against Lyme disease, not actual disease germs. In other words, the test can be negative (normal) when a child does in fact have early localized Lyme disease.

Other symptoms of early Lyme disease may accompany the rash or can occur even in the absence of the rash. This stage is called Early Disseminated disease. Within about one month from the time of the tick bite, some children with Lyme develop a rash that appears in multiple body sites all at once, not just at the site of the tick bite. Each circular lesion of rash looks like the rash described above, but usually is smaller. Additional symptoms include fever, body aches, headaches, and fatigue without other viral symptoms such as sore throat, runny nose, and cough. Some kids get one-sided facial weakness. Blood testing at this point is more likely to be positive.

The treatment of early Lyme disease is straightforward. The child takes 2-3 weeks of an antibiotic that is known to treat Lyme disease effectively such as amoxicillin or doxycycline. Your pediatrician needs to see the rash and evaluate other symptoms to make the diagnosis. Treatment prevents later

complications of the disease. Treated children fortunately do not get “chronic Lyme disease.” Once treatment is started, the rash fades over several days and other symptoms, if present, resolve. Sometimes at the beginning of treatment the child experiences chills, aches, or fever for a day or two. This reaction is normal but you should contact your child’s doctor if it persists for longer.

Later stages of Lyme disease may be treated with the same oral antibiotic as for early Lyme but for 4 weeks instead of 2-3 weeks. The most common symptom of late stage Lyme disease is arthritis (red, swollen, mildly painful joint) of a large joint such as a knee, hip, or shoulder. Some kids just develop joint swelling without pain and the arthritis can come and go.

For some manifestations, IV antibiotics are used. The longest course of treatment is 4 weeks for any stage. Again, children do not develop “chronic Lyme” disease. If symptoms persist despite adequate treatment, sometimes one more course of antibiotics is prescribed, but if symptoms continue, the diagnosis should be questioned. No advantage is shown by longer treatments. Some adults have lingering symptoms of fatigue and aches years after treatment for Lyme disease. While the cause of the symptoms is not understood, we do know that prolonged courses of antibiotics do not affect symptoms.

For kids eight years old or older, if a blacklegged tick has been attached for well over 36 hours and is clearly engorged, and if you live in an area of high rates of Lyme disease-carrying ticks, your pediatrician may in some instances choose to prescribe a one time dose of the antibiotic doxycycline to prevent Lyme disease. The study that this strategy was based on and a few other criteria that are considered in this situation are described [here](#). Your pediatrician can discuss the pros and cons of this treatment.

Bug checks and insect repellent. Protect kids with [DEET containing insect repellents](#). The Centers for Disease

Control recommends 10 to 30 percent DEET- higher percent stays on longer. Spray on clothing and exposed areas and do not apply to babies under two months of age. Grab your kids and preform daily bug checks- in particular look in crevices where ticks like to hide such as the groin, armpits, between the toes and check the hair. Ticks can be tough to spot. Dr. Lai once had a elementary school patient who had a blacklegged tick in the middle of his forehead. The mother noticed it at breakfast, tried to brush it off, thought it was a scab and sent the boy to school. Later that day the teacher called saying, "I think your son has a bug on his face."

Misinformation about this disease abounds, and self proclaimed "Lyme disease experts" play into people's fears. While pediatricians who practice in Lyme disease endemic areas are usually well versed in Lyme disease, if you feel that you need another opinion about your child's Lyme disease, the "expert" that you should consult would be a pediatric infectious disease specialist.

For a more detailed discussion of Lyme disease, look to the Center for Disease Control website: www.cdc.gov.

Julie Kardos, MD and Naline Lai, MD

©2017, 2016, 2009 Two Peds in a Pod®

Lyme Disease: What Makes it

Tic? Ticks!

As we are in the middle of Lyme disease season here in the Northeastern United states, I thought I should address Lyme disease. I have diagnosed 8 cases so far this summer, seven in my office and one at a picnic, and what struck me in each case was how relieved the parents were to find out how easy it is to treat the disease when it is diagnosed early. It is important to treat Lyme disease in the early phase because this treatment prevents later manifestations of the illness (arthritis, meningitis, etc.).



Lyme disease is spread to people by deer ticks. Any one deer tick that you pull off your child has only a 1% chance of transmitting Lyme disease, but the reason so many people get Lyme disease is that there are an awful lot of deer ticks out there.

In areas where Lyme disease is prevalent (New England and Mid-Atlantic states, upper Midwest states, and California), parents should be vigilant about searching their children's bodies daily for ticks and for the rash of early Lyme disease. Tick bites, and therefore the rash as well, especially like to show up on the head, in belt lines, groins, and axillas (armpits), but can occur anywhere. I shower my kids daily in summer time not just to wash off pool water, sunscreen, and dirt, but also for the opportunity to check them for ticks and rashes.

Most kids do get the classic rash of Lyme disease at the site of a tick bite. The rash most commonly occurs by 1-2 weeks after the tick bite and is round, flat, and typically red. It can have some central clearing. The key is that the rash

expands and becomes larger than 5cm. Untreated, it can become quite large as seen in the above photo. The rash does not itch or hurt. This finding is helpful because if you think you are seeing the primary rash of Lyme disease on your child, you can safely wait a day or two before bringing your child to his health care provider because the rash will continue to grow. The Lyme disease rash does not come and then fade in the same day. In fact, the history of a rash that enlarges over a few days is helpful in diagnosing the disease. Some kids have fever, headache, or muscle aches at the same time that the rash appears.

The second phase of Lyme disease occurs if it is not treated in the primary phase. It occurs about one month from the time of tick bite. Children develop a rash that looks like the primary rash but appears in multiple body sites all at once, not just at the site of the tick bite. Each circular lesion of rash looks like the primary rash but typically is smaller. Additional symptoms include fever, body aches, headaches, and fatigue without other viral symptoms such as sore throat, runny nose, and cough. Some kids get the fever but no rash. Some kids get one-sided facial weakness. This stage is called Early Disseminated disease and is treated similarly to the way that Early Lyme disease is treated.

If your child has primary Lyme disease (enlarging red round rash), the diagnosis is made on clinical presentation alone. **No blood work is needed** because it takes several weeks for a person's body to make antibodies to the disease, and blood work tests for antibody response. In other words, the test can be negative when a child does have early Lyme disease. Therefore, treatment begins after taking a history and performing a visual diagnosis.

The treatment of early Lyme disease is straightforward. The child takes 2-3 weeks of an antibiotic that is known to treat Lyme disease effectively such as amoxicillin or doxycycline prescribed by your child's health care provider. This

treatment prevents later complications of the disease. While the disease can progress if no treatment is undertaken, in children there is no evidence of “chronic Lyme disease” despite claims to the contrary. Once treatment is started, the rash fades over several days. Sometimes at the beginning of treatment the child experiences chills, aches, or fever for a day or two. This reaction is normal but your child’s health care provider should be contacted if it persists for longer.

If not treated early, then treatment starts when diagnosis is made during later stages of Lyme disease and may include the same oral antibiotic as for early Lyme but for 4 weeks instead of 2-3 weeks. The most common symptom of late stage Lyme disease is arthritis (red, swollen, painful joint) of a large joint such as a knee, hip, shoulder. Some kids just develop joint swelling without pain. The arthritis can come and go. This stage is prevented by early treatment but is also can be treated with antibiotics.

For some manifestations, IV antibiotics are used. The longest course of treatment is 4 weeks for any stage. Again, children do not develop “chronic Lyme” disease. If symptoms persist despite adequate treatment, sometimes one more course of antibiotics is prescribed, but if symptoms continue, the diagnosis should be questioned. No advantage is shown by longer treatments.

Misinformation about this disease abounds, and self proclaimed “Lyme disease experts” play into people’s fears. If you feel that you need another opinion about your child’s Lyme disease, the “expert” that you could consult would be a pediatric infectious disease specialist.

For a more detailed discussion of Lyme disease, I refer you to the Center for Disease Control website: www.cdc.gov.

Julie Kardos, MD with Naline Lai, MD

2009 Two Peds in a Pod, updated 2015

